

Polar Ocean Phytoplankton diversity

Investigating the biogeochemistry of the high latitudes during the period of rapid change: modelling and satellite retrievals

A. Bracher, S. N. Losa, M.A. Soppa, M. Losch, V. Pefanis, J. Oelker, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research; Institute of Environmental Physics, University of Bremen

In Short

- Polar regions
- Long time series of chlorophyll *a* for different PFTs
- Satellite retrievals and numerical modeling

The aim of this computing project is to obtain long-term (over decades) time series of ocean biogeochemical model simulations and satellite retrievals to analyse the changes in biodiversity and biogeochemical cycling observed over the last 20–30 years in the Polar regions (the Southern and Arctic Oceans) and to improve our understanding of possible interactions between the open water, sea ice, snow, ocean biogeochemistry and ecosystem and chemical composition of the Atmospheric Boundary Layer under the recently observed climate changes. Within the HLRN project, the coloured dissolved organic matter (CDOM) and phytoplankton dynamics as well as phytoplankton diversity in response to Arctic Amplification are simulated with the biogeochemical model Darwin [1,2] and REcoM [3] coupled to the Massachusetts Institute of Technology General Circulation Model (MITgcm, [4]). Satellite retrievals of chlorophyll *a* concentrations (Chla) for various phytoplankton functional types (PFTs) – diatoms, coccolithophores, cyanobacteria - are derived based on synergistic use (SynSenPFT, [5]) of multi-spectral-based [6,7] and hyper-spectral-based [8,9] phytoplankton absorption information. The SynSenPFT (Figure 1 and 2) will be further extended and updated to account for more information from current hyper-spectral (OMI) measurements and new multi-spectral Sentinel-3 data with better spatial and temporal coverage. The combined model and satellite-derived information on PFTs and CDOM absorption will be used to investigate existing relationships and feedbacks between the Arctic climate change, the ocean biogeochemistry and atmospheric oxidative capacity, which is one of the scientific tasks of the related project "Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms (AC³)" within the establishment of Transregional Collaborative Research Centre TR

172. The time series of the satellite PFT Chla retrievals and MITgcm-Darwin biogeochemical model integrations (over the period of 1991 – 2016) for the Southern Ocean are to support the DFG research "Antarctic phytoplankton in response to environmental changes studied by a synergetic approach using multi- and hyper-spectral satellite data (PhySen)" within the framework of the DFG-Priority Program 1158 "Antarctic Research".

For independent evaluation of the model and satellite retrieved PFT data we use *in situ* marker phytoplankton pigments determined with high precision liquid chromatography (HPLC) sampled and compiled by AWI "Phytooptics" team in cooperation with other researchers from AWI since 2009 onwards [10] and within the currently running HGF-project FRAM in the Arctic Ocean (Greenland Sea, Laptev Sea, Central Arctic).

The already allocated NPLs was used for sensitivity studies and model calibration that allowed to improve significantly the simulated phenology and dominance of the observed PFTs in the Southern Ocean (Figure 3) and provide recommendations for the Arctic Ocean set up.

WWW

<https://www.awi.de/ueberuns/organisation/mitarbeiter/astrid-bracher.html>

More Information

- [1] M. J. Follows, S. Dutkiewicz, S. Grant, S. W. Chisholm, Emergent biogeography of microbial communities in a model Ocean, *Science* **315**, 1843 – 1846 (2007). doi: 10.1126/science.1138544
- [2] S. Dutkiewicz, A. E. Hickman, O. Jahn, W. W. Gregg, C. B. Mouw, M. J. Follows, Capturing optically important constituents and properties in a marine biogeochemical and ecosystem model, *Biogeosciences* **12**, 4447 – 4481 (2015). doi:10.5194/bg-12-4447-2015
- [3] M. H. Taylor, M. Losch, A. Bracher, On the drivers of phytoplankton blooms in the Antarctic seasonal ice zone: a modelling approach, *J. Geophys. Res. – Oceans* **188**, 63 – 75 (2013). doi:10.1029/2012JC008418
- [4] MITgcm Group, MITgcm Manual, Online documentation, *MIT/EAPS, Cambridge, MA 02139, USA* (2012).

- [5] S. N. Losa, M. A. Soppa, T. Dinter, A. Wolanin, R. J. W. Brewin, A. Bricaud, J. Oelker, I. Peeken, B. Gentili, V. V. Rozanov, A. Bracher, Synergistic exploitation of hyper- and multispectral Sentinel measurements to determine Phytoplankton Functional Types (SynSenPFT), *Frontiers in Marine Science* **4**, 203 (2017). doi:10.3389/fmars.2017.00258; doi:10.3389/fmars.2017.00203
- [6] M. A. Soppa, T. Hirata, B. Silva, T. Dinter, I. Peeken, S. Wiegmann, A. Bracher, Global Retrieval of Diatom Abundance Based on Phytoplankton Pigments and Satellite Data, *Remote Sensing* **6**(19), 10089 – 10106 (2014). doi:10.3390/rs61010089
- [7] M. A. Soppa, I. Peeken, A. Bracher, Global chlorophyll "a" concentrations for diatoms, haptophytes and prokaryotes obtained with the Diagnostic Pigment Analysis of HPLC data compiled from several databases and individual cruises (2017). doi:10.1594/PAN-GAEA.875879
- [8] A. Bracher, M. Vountas, T. Dinter, J. P. Burrows, R. Röttgers, I. Peeken, Quantitative observation of cyanobacteria and diatoms from space using PhytoDOAS on SCIAMACHY data, *Biogeosciences* **6**, 751 – 764 (2009). doi:10.5194/bg-6-751-2009
- [9] A. Sadeghi, T. Dinter, M. Vountas, B. B. Taylor, M. A. Soppa, I. Peeken, A. Bracher, Improvements to the PhytoDOAS method for identification of coccolithophores using hyperspectral satellite data, *Ocean Sciences* **8**, 1055 – 1070 (2012). doi:10.5194/bg-9-2127-2012
- [10] E. M. Nöthig, A. Bracher, A. Engel, K. Metfies, B. Niehoff, I. Peeken et al., Plankton ecology and biogeochemistry in a changing arctic ocean – overview of long-term summer measurements in the Fram strait variability of chlorophyll a distribution in the Fram Strait, Greenland sea and central Arctic Ocean, *Polar Research* **34** (2015). doi:10.3402/polar.v34.23349

Funding

DFG-Priority Program 1158 "Antarctic Research"; DFG Transregional Collaborative Research Centre TR 172; Helmholtz Climate Initiative REKLIM (Regional Climate Change), a joint research project of the Helmholtz Association of German Research Centres (HGF)

SynSenPFT algorithm




Figure 1: Diagram exemplifying the SynSenPFT algorithm that updates the PFT information from each OC-PFT sub-pixel (x_b) within a PhytoDOAS pixel given the PhytoDOAS Chla (y) and a priori error statistics introduced in K , which results in SynSenPFT Chla (x_a) at each OC-PFT sub-pixel.




Figure 2: SynSenPFT retrievals of coccolithophores Chla.




Figure 3: Simulated spatial distribution of the dominant phytoplankton groups in the Southern Ocean.