


# FOCI - a new climate model

## Coupled climate modeling with a high resolution ocean

**J. Harlass and A. Biastoch**, GEOMAR Helmholtz Centre for Ocean Research Kiel

### In Short

- New state-of-the-art coupled climate model with a high-resolution nested ocean at GEOMAR
- Role of an active atmosphere on the effect of mesoscale ocean eddies on decadal variability of the large-scale oceanic circulation
- First study with a high-resolution nest in the Indian/South Atlantic Ocean



**Figure 1:** Schematic of the new model system at GEOMAR: Flexible Ocean and Climate Infrastructure - FOCI. Some components are optional.

Mesoscale eddies are one of the most important features for the dynamic of major current systems. They also play a crucial role for the decadal variability of the global overturning circulation [1]. Yet, most ocean general circulation models still have too coarse resolution to explicitly resolve such mesoscale eddies; even most modern climate models rely on parameterizations of the eddy effect [2]. By increasing spatial resolution over the global domain to  $1/10^\circ$  and finer, the mesoscale can be captured at the expense of extreme computational costs and hence comparably short simulations. Our approach is to increase ocean resolution in specific and outstanding areas of interest: so-called regional nesting (AGRIF [3]).

At GEOMAR we successfully implemented high resolution ocean nesting into a fully coupled atmosphere-ocean climate model, depicted schematically in Fig. 1. With the Flexible Ocean and Climate Infrastructure (FOCI) we can for the first time study ocean-atmosphere feedbacks in an eddying ocean and start to assess their importance for decadal climate variability.

A great advantage of the ocean nesting is its lower computational costs, as compared to globally high-resolved ocean, and hence the ability to perform series of multi-decadal simulations.

The global ocean, based on current NEMO 3.6 code, has a  $1/2^\circ$  spatial resolution (ORCA05), while in the nested area resolution increases to  $1/10^\circ$ . A first ocean nest has been configured in the Indian and South Atlantic Ocean *INALT10* ( $70^\circ\text{W}-70^\circ\text{E}$ ,  $50^\circ\text{S}-10^\circ\text{N}$ , detailed description of precursor nest *INALT01* in [4]). In the mean time, a second independent nest has been configured in the North Atlantic (*VIKING10*, project shk00029) focusing on the sensitivity of the climate system to freshwater input into the sub-polar North Atlantic (Greenland

melting).


The atmosphere model ECHAM (version 6.3) has a horizontal resolution of T63 ( $\sim 1.875^\circ$ ) and 95 vertical levels with a top level at 0.01hPa (80km). JSBACH deals with all land processes and does also provide river runoff for the ocean.

Exchange of momentum, sea-ice properties and heat and freshwater fluxes between both components is performed by the OASIS3-MCT coupler. Technically, the atmosphere receives the interface fields from the global ocean base grid, but a 2-way-nesting technique ensures, that information from inside the high resolution nest transfer to the coarser base model grid.

In this project our main focus lies on the impact of mesoscale ocean eddies on decadal variability of the large-scale oceanic circulation and in particular the role of an active atmosphere. With the new coupled model and the *INALT10* nest we want to explore the role of mesoscale eddies for atmosphere-ocean interactions.

For our studies, the global coupled (non-nested) model is integrated under pre-industrial climate conditions for 1000 year to spin-up the climate system. The nested *INALT10* experiment is then restarted at year 1000 and integrated for another 100 years. The *INALT10* nested configuration runs on 51 nodes and takes less than 2 hours, while the non-nested version needs only 43 minutes on 40 nodes, to simulate one model year.

Resolving smaller ocean scales reduces a prominent bias in current state-of-the-art climate models [7], the sea surface temperature (SST) warm bias in the tropical Atlantic, significantly, Fig.



**Figure 2:** Evolution of Sea Surface Temperature (SST) bias in boreal summer, for a) non-nested, b) INALT10 nest and c) difference.

2. The SST bias in the non-nested FOCI closely resembles the ensemble mean of the climate models participating in CMIP5, in terms of pattern size and magnitude. The cooling in the equatorial and Benguela upwelling region by about 2K, Fig. 2c, is also reflected in improved precipitation and wind pattern along the equatorial Atlantic (not shown). We would like to analyze the role of an active atmosphere on the impact of Agulhas mesoscale on the Atlantic Meridional Overturning Circulation (AMOC) as just high-resolution models reproduce the observed salt import into the Atlantic that impacts stability of the Gulf Stream system [5]. In the performed experiments so far, AMOC at 26°N has a strength of about 18Sv in non-nested and nested FOCI. A secondary spin-up time of 50-60 years became apparent in the nested coupled experiments, that needs to be taken into account in future experiments.

Our research will further focus on ocean resolution dependence for e.g. tropical Atlantic warming [6] or Benguela upwelling in the context of coupled atmosphere-ocean feedbacks.

- [4] J.V. Durgadoo, B.R. Loveday, C.J.C Reason, et al., Agulhas Leakage Predominantly Responds to the Southern Hemisphere Westerlies. *J. Phys. Oceanogr.* **43**, (2013). doi: 10.1175/JPO-D-13-047.1
- [5] J. Deshayes, A.-M. Treguier, B. Barnier, et al., Oceanic hindcast simulations at high resolution suggest that the Atlantic MOC is bistable. *Geophys. Res. Lett.* **40**, (2013). doi: 10.1002/grl.50534
- [6] J. Luebbecke, J.V. Durgadoo, A. Biastoch, Contribution of Increased Agulhas Leakage to Tropical Atlantic Warming. *J. Clim.* **28**, (2015). doi:10.1175/JCLI-D-15-0258.1
- [7] I. Richter, S.P. Xie, S. Behera, et al., Equatorial Atlantic variability and its relation to mean state biases in CMIP. *Clim. Dyn.* **42**, (2012). doi: 10.1007/s00382-012-1624-5

### Funding

Helmholtz Association - Earth System Modeling

### WWW

<http://www.geomar.de/go/ocean-models>

### More Information

- [1] A. Biastoch, J.V. Durgadoo et al., Atlantic multi-decadal oscillation covaries with Agulhas leakage. *Nat. Commun.* **6**, (2015). doi: 10.1038/ncomms10082
- [2] R. Hallberg, Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects. *Ocean Model.* **72**, (2013). doi: 10.1016/j.ocemod.2013.08.007
- [3] L. Debreu, E. Blayo, Two-way embedding algorithms: a review. *Ocean Dyn.* **58**, (2008). doi:10.1007/s10236-008-0150-9